

On new modular sequence space defined over 2-normed spaces

Gülcan ATICI TURAN*

Department of Mathematics, University of Muş Alparslan, Muş, Turkey

Geliş Tarihi (Received Date): 06.07.2019

Kabul Tarihi (Accepted Date): 17.09.2019

Abstract

In this paper, a new sequence space $F(\|\cdot, \cdot\|, \mathcal{M}, p, u)$ is defined by using a sequence of Orlicz functions in 2-normed spaces. Some various properties and some inclusions are also examined on this space.

Keywords: Orlicz function, sequence spaces, 2-norm, paranormed spaces.

2-normlu uzaylarda tanımlı yeni modular dizi uzayı

Öz

Bu çalışmada, 2-normlu uzaylarda Orlicz fonksiyonlarının bir dizisi kullanılarak $F(\|\cdot, \cdot\|, \mathcal{M}, p, u)$ yeni dizi uzayı tanımlanmıştır. Ayrıca bu uzayın bazı özellikleri ve bazı kapsama bağıntıları incelenmiştir.

Anahtar Kelimeler: Orlicz fonksiyon, dizi uzayları, 2-norm, paranormlu uzaylar.

1. Introduction

The concept of 2-normed spaces was introduced and studied by Siegfried Gähler, a German Mathematician who worked at German Academy of Science, Berlin, in a series of paper in German language published in Mathematische Nachrichten, see for example references [1,2]. This notion which is nothing but a two dimensional analogue of a normed space got the attention of a wider audience after the publication of a paper by Albert George, White Jr. of USA in 1969 entitled 2-Banach spaces [3]. In the same year Gähler published another paper on this theme in the same journal [1]. A.H. Siddiqi

* Gülcan ATICI TURAN, gatici23@hotmail.com, <https://orcid.org/0000-0002-1009-6072>

delivered a series of lectures on this theme in various conferences in India and Iran. His joint paper with Gähler et al. [4] of 1975 also provide valuable results related to the theme of this paper. Results up to 1977 were summarized in the survey paper by Siddiqi [5].

An Orlicz function $M : [0, \infty) \rightarrow [0, \infty)$ is a function, which is continuous, nondecreasing and convex such that $M(0) = 0, M(x) > 0$ for $x > 0$ and $M(x) \rightarrow \infty$ as $x \rightarrow \infty$.

Note that for M is an Orlicz function, we have $M(\lambda x) \leq \lambda M(x)$ where $0 \leq \lambda \leq 1$
 ℓ_M sequence space defined as following:

$$\ell_M = \left\{ x \in w : \sum_{k=1}^{\infty} \left(M \left(\frac{|x_k|}{\rho} \right) \right) < \infty, \text{ for some } \rho > 0 \right\} [6].$$

Let X be a real linear space and $\|.,.\|$ is defined a real valued mapping on $X \times X$. For $x, y, z \in X$ and $\lambda \in \mathbb{R}$, the function $\|.,.\|$, which satisfies the following conditions is called 2-norm and the pair $(X, \|.,.\|)$ is called a linear 2-normed space or shortly 2-normed space. $\|.,.\|$ is a non-negative function.

- (N₁) $\|x, y\| = 0$ if and only if x and y are linearly dependent;
- (N₂) $\|x, y\| = \|y, x\|$;
- (N₃) $\|\lambda x, y\| = |\lambda| \|x, y\|, \lambda \in \mathbb{R}$;
- (N₄) $\|x, y + z\| \leq \|x, y\| + \|x, z\|$.

$(X, \|.,.\|)$ is a 2-Banach space if every Cauchy sequence in X is convergent to some x in X [7].

Let X be a linear metric space. A function $g: X \rightarrow \mathbb{R}$ is called paranorm, if

- (i) $g(x) \geq 0$, for all $x \in X$
- (ii) $g(-x) = g(x)$, for all $x \in X$
- (iii) $g(x + y) \leq g(x) + g(y)$, for all $x, y \in X$
- (iv) if (μ_n) is a sequence of scalars with $\mu_n \rightarrow \mu$ as $n \rightarrow \infty$ and (x_n) is a sequence of vectors with $g(x_n - x) \rightarrow 0$ as $n \rightarrow \infty$, then $g(\mu_n x_n - \mu x) \rightarrow 0$ as $n \rightarrow \infty$ [8].

A scalar valued paranormed sequence space (F, g_F) , where g_F is a paranorm on F is called monotone paranormed space if $x = (x_k), y = (y_k) \in F$ and $|x_k| \leq |y_k|$ for all k implies $g_F(x) \leq g_F(y)$ [8].

Definition 1.1. Let X be a sequence space.

- (i) If $y = (y_k) \in X$ whenever $|y_i| \leq |x_i|, i \geq 1$ for some $x = (x_k) \in X$, then X is called solid (or normal).
- (ii) If $(x_k) \in X$ implies $(X_{\pi(k)}) \in X$ such that $\pi(k)$ is a permutation of \mathbb{N} , then X is called symmetric [9].

U is showed as the set of all real sequences $u = (u_k)$, where $u_k > 0$ for all $k \in \mathbb{N}$.

Throughout this study the following inequality will be used. Let $p = (p_k)$ be a positive sequence of real numbers with $0 < p_k \leq \sup p_k = H, D = \max(1, 2^{H-1})$, then for all $a_k, b_k \in \mathbb{C}$, we have

$$|a_k + b_k|^{p_k} \leq D\{|a_k|^{p_k} + |b_k|^{p_k}\}. \tag{1}$$

2. Main results

Let (F, g_F) be a normal paranormed sequence space with paranorm g_F which satisfies the following properties:

- (i) g_F is a monotone paranorm;
- (ii) coordinatewise convergence implies convergence in paranorm g_F , which implies that for each $(X^n) = (X_k^n) \in F, n, k \in \mathbb{N}, X_k^n \rightarrow 0$ as $n \rightarrow \infty$ (for each k) $\Rightarrow g_F(X^n) \rightarrow 0$ as $n \rightarrow \infty$ [10].

Let $(N, \|\cdot, \cdot\|)$ be a 2-normed space and $\mathcal{M} = (M_k)$ be a sequence of Orlicz functions. Further, let $p = (p_k)$ be a bounded sequence of strictly positive real numbers. We define the set:

$$F(\|\cdot, \cdot\|, \mathcal{M}, p, u) = \left\{ X = (X_k) : X_k \in N, \left(u_k \left[M_k \left(\frac{\|X_k, Z\|}{\rho} \right) \right]^{p_k} \right) \in F, \text{ for some } \rho > 0 \right\}$$

for every $Z \in N$.

For $p_k = 1$ for all $k \in \mathbb{N}$, we write this space as $F(\|\cdot, \cdot\|, \mathcal{M}, u)$.

Theorem 2.1. If $\mathcal{M} = (M_k)$ is a sequence of Orlicz functions then $F(\|\cdot, \cdot\|, \mathcal{M}, p, u)$ is a linear space.

Proof. Let $X = (X_k), Y = (Y_k) \in F(\|\cdot, \cdot\|, \mathcal{M}, p, u)$ and $a, b \in \mathbb{R}$, thus there are some positive numbers ρ_1 and ρ_2 such that

$$\left(u_k \left[M_k \left(\frac{\|X_k, Z\|}{\rho_1} \right) \right]^{p_k} \right) \in F \quad \text{and} \quad \left(u_k \left[M_k \left(\frac{\|Y_k, Z\|}{\rho_2} \right) \right]^{p_k} \right) \in F$$

for every $Z \in N$. Define $\rho = \max\{2|a|\rho_1, 2|b|\rho_2\}$. Because of the definition of the Orlicz function, we can write

$$\begin{aligned} u_k \left[M_k \left(\frac{\|aX_k + bY_k, Z\|}{\rho} \right) \right]^{p_k} &\leq u_k \left[M_k \left(\frac{\|aX_k, Z\| + \|bY_k, Z\|}{\rho} \right) \right]^{p_k} \\ &< u_k \left[M_k \left(\frac{\|X_k, Z\|}{\rho_1} \right) + M_k \left(\frac{\|Y_k, Z\|}{\rho_2} \right) \right]^{p_k} \\ &\leq Du_k \left[M_k \left(\frac{\|X_k, Z\|}{\rho_1} \right) \right]^{p_k} + Du_k \left[M_k \left(\frac{\|Y_k, Z\|}{\rho_2} \right) \right]^{p_k} \in F, \end{aligned}$$

such that $D = \max\{1, 2^{H-1}\}$. Therefore $aX + bY \in F(\|\cdot, \cdot\|, \mathcal{M}, p, u)$. Hence $F(\|\cdot, \cdot\|, \mathcal{M}, p, u)$ is a linear space.

Theorem 2.2. For any sequence $\mathcal{M} = (M_k)$ of Orlicz function, $F(\|\cdot, \cdot\|, \mathcal{M}, p, u)$ is a paranormed space with

$$g_T(X) = \inf \left\{ \rho^{\frac{p_k}{T}} > 0 : \left[g_F \left(u_k \left[M_k \left(\frac{\|X_k, Z\|}{\rho} \right) \right]^{p_k} \right) \right]^{\frac{1}{T}} \leq 1, k = 1, 2, \dots \right\} \quad (2)$$

such that $T = \max(1, H)$, $H = \sup_k p_k < \infty$ and $\inf p_k > 0$ and for $Z \in N$.

Proof. It is easy to prove that $g_T(\theta) = 0$ and $g_T(-X) = g_T(X)$. Since g_F is monotone and when $a = b = 1$ is taken in the proof of Theorem 2.1, we write $g_T(X + Y) \leq g_T(X) + g_T(Y)$ for $X = (X_k), Y = (Y_k) \in F(\|\cdot, \cdot\|, \mathcal{M}, p, u)$.

Let $\lambda \neq 0$ be any complex number. Because of the continuity of the scalar multiplication, we can write

$$\begin{aligned} g_T(\lambda X) &= \inf \left\{ \rho^{\frac{p_k}{T}} > 0 : \left[g_F \left(u_k \left[M_k \left(\frac{\|\lambda X_k, Z\|}{\rho} \right) \right]^{p_k} \right) \right]^{\frac{1}{T}} \leq 1, k = 1, 2, \dots \right\} \\ &= \inf \left\{ (|\lambda|r)^{\frac{p_k}{T}} > 0 : \left[g_F \left(u_k \left[M_k \left(\frac{\|X_k, Z\|}{r} \right) \right]^{p_k} \right) \right]^{\frac{1}{T}} \leq 1, k = 1, 2, \dots \right\} \end{aligned}$$

where $r = \rho/|\lambda|$.

Since $|\lambda|^{p_k} \leq \max(1, |\lambda|^H)$. We have $|\lambda|^{\frac{p_k}{T}} \leq (\max(1, |\lambda|^H))^{\frac{1}{T}}$. Thus $g_T(\lambda X)$ converges to zero when $g_T(X)$ converges to zero in $F(\|\cdot, \cdot\|, \mathcal{M}, p, u)$.

Let $X = (X_k) \in F(\|\cdot, \cdot\|, \mathcal{M}, p, u)$ and assume that $\lambda_n \rightarrow 0$ as $n \rightarrow \infty$. Let $\varepsilon > 0$ and K be a positive integer. Then we can write

$$g_F \left(u_k \left[M_k \left(\frac{\|X_k, Z\|}{\rho} \right) \right]^{p_k} \right) < \left(\frac{\varepsilon}{2} \right)^T$$

every some $\rho > 0$ and for $k > K$ such that $k \in N$,

$$\left[g_F \left(u_k \left[M_k \left(\frac{\|X_k, Z\|}{\rho} \right) \right]^{p_k} \right) \right]^{\frac{1}{T}} \leq \frac{\varepsilon}{2}.$$

Let $0 < |\lambda| < 1$. Because of the definition of the Orlicz function and by the condition (iii) of 2-norm, we have

$$\begin{aligned} g_F \left(u_k \left[M_k \left(\frac{\|\lambda X_k, Z\|}{\rho} \right) \right]^{p_k} \right) &= g_F \left(u_k \left[M_k \left(|\lambda| \frac{\|X_k, Z\|}{\rho} \right) \right]^{p_k} \right) \\ &< g_F \left(u_k \left[|\lambda| M_k \left(\frac{\|X_k, Z\|}{\rho} \right) \right]^{p_k} \right) \end{aligned}$$

$$\begin{aligned} &< g_F \left(u_k \left[M_k \left(\frac{\|X_k, Z\|}{\rho} \right) \right]^{p_k} \right) \\ &< \left(\frac{\varepsilon}{2} \right)^T \end{aligned}$$

for $k > K$. Since M is continuous everywhere in $[0, \infty)$ and by the definition of g_F , it follows that for $k \leq K$

$$\varphi(t) = g_F \left(u_k \left[M_k \left(\frac{\|tX_k, Z\|}{\rho} \right) \right]^{p_k} \right)$$

is continuous at 0. Therefore $|\varphi(t)| < \frac{\varepsilon}{2}$ for $0 < t < \delta$ such that $0 < \delta < 1$. Let L be any integer such that $|\lambda_n| < \delta$ for $n > L$, then

$$\left[g_F \left(u_k \left[M_k \left(\frac{\|\lambda_n X_k, Z\|}{\rho} \right) \right]^{p_k} \right) \right]^{\frac{1}{T}} < \frac{\varepsilon}{2}$$

for $n > L$ and $k \leq K$. Therefore

$$\left[g_F \left(u_k \left[M_k \left(\frac{\|\lambda_n X_k, Z\|}{\rho} \right) \right]^{p_k} \right) \right]^{\frac{1}{T}} < \varepsilon$$

for $n > L$ and for all k . So $\lambda_n X \rightarrow \theta$ as $n \rightarrow \infty$. This completes the proof of the theorem.

Theorem 2.3. Let $(N, \|\cdot, \cdot\|)$ be a 2-Banach space, then the space $F(\|\cdot, \cdot\|, \mathcal{M}, p, u)$ is a complete paranormed space with $g_T(X)$, where F is a K -space.

Proof. The proof is routine verification by using standard arguments and therefore omitted.

Theorem 2.4. If F is a K -space, then $F(\|\cdot, \cdot\|, \mathcal{M}, p, u)$ is a K -space.

Proof. Let us define a mapping $\tau_n: F(\|\cdot, \cdot\|, \mathcal{M}, p, u) \rightarrow N$ by $\tau_n(X) = X_n, \forall n \in \mathbb{N}$. Our aim is to show τ_n is continuous.

Let (X^m) be a sequence in $F(\|\cdot, \cdot\|, \mathcal{M}, p, u)$ such that $X^m \xrightarrow{g} 0$ as $m \rightarrow \infty$. Then for some suitable choice of $\rho > 0$,

$$\left[g_F \left(u_k \left[M_k \left(\frac{\|X_k^m, Z\|}{\rho} \right) \right]^{p_k} \right) \right]^{\frac{1}{T}} \rightarrow 0$$

as $m \rightarrow \infty$. Since F is a K -space, this implies that for each k and as m tending to ∞ ,

$$u_k \left[M_k \left(\frac{\|X_k^m, Z\|}{\rho} \right) \right]^{p_k} \rightarrow 0$$

for some $\rho > 0$. Since M_k be a sequence of Orlicz functions, it follows that $\|X_k^m, Z\| \rightarrow 0$ as $m \rightarrow \infty$. Consequently, $X^m \rightarrow 0$ in N .

Theorem 2.5. Let \mathcal{M} and \mathcal{T} be two sequence of Orlicz functions. Then

$$F(\|\cdot, \cdot\|, \mathcal{M}, p, u) \cap F(\|\cdot, \cdot\|, \mathcal{T}, p, u) \subseteq F(\|\cdot, \cdot\|, \mathcal{M} + \mathcal{T}, p, u)$$

where F is a normal sequence space.

Proof. Let $X = (X_k) \in F(\|\cdot, \cdot\|, \mathcal{M}, p, u) \cap F(\|\cdot, \cdot\|, \mathcal{T}, p, u)$. Then we can choose $\rho_1, \rho_2 > 0$ such that

$$\left(u_k \left[M_k \left(\frac{\|X_k, Z\|}{\rho_1} \right) \right]^{p_k} \right) \in F \text{ and } \left(u_k \left[T_k \left(\frac{\|X_k, Z\|}{\rho_2} \right) \right]^{p_k} \right) \in F.$$

Define $\rho = \max\{\rho_1, \rho_2\}$. We can write

$$u_k \left[(M_k + T_k) \left(\frac{\|X_k, Z\|}{\rho} \right) \right]^{p_k} \leq u_k D \left\{ \left[M_k \left(\frac{\|X_k, Z\|}{\rho_1} \right) \right]^{p_k} + \left[T_k \left(\frac{\|X_k, Z\|}{\rho_2} \right) \right]^{p_k} \right\} \in F,$$

where $D = \max\{1, 2^{H-1}\}$. Since F is normal, $X \in F(\|\cdot, \cdot\|, \mathcal{M} + \mathcal{T}, p, u)$.

Theorem 2.6. Let $\mathcal{M} = (M_k)$ be a sequence of Orlicz functions. Then $c_0(\|\cdot, \cdot\|, \mathcal{M}, p, u) \subset c(\|\cdot, \cdot\|, \mathcal{M}, p, u) \subset \ell_\infty(\|\cdot, \cdot\|, \mathcal{M}, p, u)$.

Proof. It is obvious that $c_0(\|\cdot, \cdot\|, \mathcal{M}, p, u) \subset c(\|\cdot, \cdot\|, \mathcal{M}, p, u)$. The second inclusion follows from the following inequality. Let $X = (X_k) \in c(\|\cdot, \cdot\|, \mathcal{M}, p, u)$ and for some $\rho = 2\mu > 0$, we obtain

$$u_k \left[M_k \left(\frac{\|X_k, Z\|}{\rho} \right) \right]^{p_k} \leq u_k D \left[M_k \left(\frac{\|X_k - L, Z\|}{\mu} \right) \right]^{p_k} + u_k D \max \left\{ 1, \left[M_k \left(\frac{\|L, Z\|}{\mu} \right) \right]^{p_k} \right\}.$$

Thus $X = (X_k) \in \ell_\infty(\|\cdot, \cdot\|, \mathcal{M}, p, u)$.

Theorem 2.7. Let $\mathcal{M} = (M_k)$ be a sequence of Orlicz functions. Then

- (i) If $0 < \inf p_k \leq p_k \leq 1$, then $c_0(\|\cdot, \cdot\|, \mathcal{M}, u) \subset c_0(\|\cdot, \cdot\|, \mathcal{M}, p, u)$;
- (ii) If $1 \leq p_k \leq \sup p_k < \infty$, then $c_0(\|\cdot, \cdot\|, \mathcal{M}, p, u) \subset c_0(\|\cdot, \cdot\|, \mathcal{M}, u)$.

Proof. (i) Let $X = (X_k) \in c_0(\|\cdot, \cdot\|, \mathcal{M}, u)$. Since $0 < \inf p_k \leq p_k \leq 1$, then we have

$$\left[M_k \left(\frac{\|X_k, Z\|}{\rho} \right) \right]^{p_k} \leq M_k \left(\frac{\|X_k, Z\|}{\rho} \right).$$

Therefore $X = (X_k) \in c_0(\|\cdot, \cdot\|, \mathcal{M}, p, u)$.

(ii) Let $1 \leq p_k \leq \sup p_k < \infty$ and $X = (X_k) \in c_0(\|\cdot, \cdot\|, \mathcal{M}, p, u)$. Then for each $0 < \varepsilon < 1$ there is a positive integer L such that

$$u_k \left[M_k \left(\frac{\|X_k, Z\|}{\rho} \right) \right]^{p_k} \leq \varepsilon < 1, \quad \forall k \geq L.$$

Since $1 \leq p_k \leq \sup p_k < \infty$, then we have

$$u_k \left[M_k \left(\frac{\|X_k, Z\|}{\rho} \right) \right] \leq u_k \left[M_k \left(\frac{\|X_k, Z\|}{\rho} \right) \right]^{p_k}.$$

Therefore $X = (X_k) \in c_0(\|\cdot, \cdot\|, \mathcal{M}, u)$. This completes the proof of the theorem.

Theorem 2.8. The space $F(\|\cdot, \cdot\|, \mathcal{M}, p, u)$ is both solid(normal) and symmetric.

Proof. The proof is similar to [10].

References

- [1] Gähler, S., Über 2-Banach räume, **Mathematische Nachrichten**, 42, 335-347, (1969).
- [2] Diminnie, C., Gähler, S. and White Jr. A.G., 2-inner product spaces, **Demonstrario Mathematica**, 6, 525-536, (1973).
- [3] White Jr. A.G., 2-Banach spaces, **Mathematische Nachrichten**, 42, 43-60, (1969).
- [4] Gähler, S., Siddiqi, A.H. and Gupta, S.C., Contributions to non-archimedean functional analysis, **Mathematische Nachrichten**, 69, 162-171, (1975).
- [5] Siddiqi, A.H., 2-normed spaces, **The Aligarh Bulletin of Mathematics**, 53-70, (1980).
- [6] Lindenstrauss, J., Tzafriri, L. On Orlicz sequence spaces, **Israel Journal of Mathematics**, 101, 379-390, (1971).
- [7] Gähler, S., Lineare 2-normierte Räume, **Mathematische Nachrichten**, 28, 1-43, (1965).
- [8] Maddox, I. J., **Elements of functional analysis**, Cambridge University Press, Cambridge, (1970).
- [9] Kamthan, P.K., Gupta, M., **Sequence spaces and series**, Marcel Dekker, New York, (1981).
- [10] Dutta, H., Kılıçman, A., Altun, Ö., Topological properties of some sequences defined over 2-normed spaces, **SpringerPlus**, 5, 2-16, (2016).
- [11] Dutta, H., Some statistically convergent difference sequence spaces defined over real 2-normed linear spaces, **Applied Sciences**, 12, 37-47, (2010).
- [12] Bektaş, Ç.A., Altin, Y., The sequence space $l_M(p, q, s)$ on seminormed spaces, **Indian Journal of Pure Applied Mathematics**, 34(4), 529-534, (2003).
- [13] Rao, M.M., Ren, Z.D., **Theory on Orlicz spaces**, Marcel Dekker, New York, (1991).
- [14] Gürdal, M. and Şahiner, A., Ideal Convergence in n-normed spaces and some new sequence spaces via n-norm, **Journal of Fundamental Sciences**, 4(1), 233-244, (2008).
- [15] Şahiner, A., Gürdal, M., Saltan, S. and Gunawan, H., Ideal convergence in 2-normed spaces, **Taiwanese Journal of Mathematics**, 11(5), 1477-1484, (2007).
- [16] Şahiner, A. and Gürdal, M., New sequence spaces in n-normed spaces with respect to an Orlicz function, **The Aligarh Bulletin of Mathematics**, 27(1), 53-58, (2008).